

Farmington Valley Health District COVID-19 Decision Tree for PreK-12 Students*

Updated 01-05-2022

If you have had a **known exposure** to someone with COVID-19:

If you are fully vaccinated against COVID-19:

2 weeks after second dose of Pfizer or Moderna vaccine or 2 weeks after dose of J & J vaccine

Then:

1. Continue with in-person learning
2. Wear a well-fitting mask around others for 10 days from last day of exposure
3. Test on day 5 from from the last day of exposure, if possible
4. **If you develop symptoms, stay home, isolate, and get a test as soon as possible**

If you are not fully vaccinated against COVID-19:

One dose of Pfizer or Moderna vaccine or within 2 weeks of final dose of any vaccine

OR

If you are unvaccinated:

Then:


1. Quarantine at home for 5 days from last day of exposure. You may return to school on day 6 if you remain asymptomatic.
2. Wear a well-fitting mask around others for 5 additional days
3. Test on day 5 from the last day of exposure, if possible
4. **If you develop symptoms, stay home, isolate, and get a test as soon as possible**

If you develop one or more symptoms, regardless of vaccination status: **STAY HOME AND GET TESTED**

Symptoms of COVID-19:

- Fever or chills
- Cough
- Shortness of breath or difficulty breathing
- Fatigue
- Muscle or body aches
- Headache
- Sore throat
- Loss of taste or smell
- Congestion or runny nose
- Nausea or vomiting
- Diarrhea

At this time, any type of test, including a home test, is acceptable.



If Positive:

1. Isolate for at least 5 days from symptom onset or test date if asymptomatic
2. After 5 days, you may return to school when fever-free for at least 24 hours and other symptoms are significantly improved
3. Continue to wear a well-fitting mask while around others for 5 additional days, especially while at home and at school

If Negative:

May return to school per school's protocols **and** when fever-free for at least 24 hours and other symptoms are significantly improved.

Note: those who have symptoms but test negative on a rapid test are recommended to seek PCR testing to confirm if possible.